

Angiographic Pitfall: Duplicated Tapered A1 Segment of the Anterior Cerebral Artery Mimicking an Anterior Communicating Artery Aneurysm

A.G. WEIL¹, M.W. BOJANOWSKI¹, F. SCHOLTES^{1,3}, T.E. DARSAUT², F. SIGNORELLI^{1,4}, A. WEILL²

¹ Department of Surgery, Division of Neurosurgery, Department of Radiology; ² Interventional Neuroradiology Unit, Centre Hospitalier de l'Université de Montréal (CHUM), Hôpital Notre-Dame; Montreal, QC, Canada

³ Department of Neurosurgery, Centre Hospitalier Universitaire, University of Liège; Liège, Belgium

⁴ Department of Neurosurgery, University Magna Graecia; Catanzaro, Italy

Key words: A1 duplication, A1 fenestration, aneurysm, subarachnoid hemorrhage, anterior communicating artery

Summary

We describe a misleading case of a partially occluded A1 segment duplication that mimicked an ACoA aneurysm on computed tomography angiography and conventional angiography and led to surgical intervention. The location of such an anomaly at the ACoA on the side of least hemodynamic stress may provide a clue to recognizing this variant.

Introduction

Duplication of the A1 segment of the anterior cerebral artery (ACA) occurs in up to 4% of normal subjects¹⁻⁵. Its incidence may double in patients harboring anterior communicating artery (ACoA) aneurysms^{1,4,6}. This variation is frequently undetected angiographically but discovered during surgery^{1,4,6}. We report a misleading case where a partially occluded A1 segment duplication mimicked an ACoA aneurysm on computed tomography angiography (CTA) and conventional angiography.

Case Report

A 70-year-old man with a medical history of hypertension, dyslipidemia, coronary artery disease, and a right middle cerebral artery

(MCA)-territory stroke from a known right internal carotid artery (ICA) occlusion from which he made a full recovery presented with sudden onset headache, nausea, and nuchal rigidity. Computed tomography (CT) showed subarachnoid haemorrhage in the basal cisterns and in the interhemispheric and sylvian fissures. In addition to the complete occlusion of the right ICA, 3D-CTA revealed an image highly suggestive of an aneurysm at the ACoA adjacent to the right A1-A2 junction (Figure 1). The right anterior circulation was supplied entirely by the left A1 through the ACoA. Conventional four-vessel angiography confirmed these findings (Figure 2). Endovascular treatment of the aneurysm was deemed unsafe because the right anterior circulation was dependent on the ACoA and surgical clipping was elected. Through a left pterional approach, the anomaly suspected to be an ACoA aneurysm was found to correspond to a tapered duplication of the A1 segment of the right ACA (Figure 3). There was no evidence of aneurysm at the ACoA complex. No other aneurysm or source of bleeding could be identified despite optimal operative exposure and inspection of the entire anterior circulation bilaterally up to the left MCA and right ICA bifurcation. The majority of blood was located around the left ICA. The patient went on to develop clinical and radiographic vasospasm, which was treated with medical therapy. Repeat good quality an-




Figure 1 3D CT-angiography revealing what was thought to be an aneurysm originating from the right side of the ACoA adjacent to the R A1-A2 junction. There is complete occlusion of the right ICA and the right anterior circulation is supplied entirely by the left ICA.




Figure 2 Conventional angiography of the left internal carotid artery (A, B) with 3D reconstructions (C) showing the supposed aneurysm originating from the ACommA complex (arrows).




Figure 3 Intraoperative photograph through a left pterional approach showing the duplicated A1 segment of the right ACA. The proximal A1 segment (arrow) is known to be occluded on angiogram. Circulation through the A2 segment (black arrow) arises from the left A1 segment (black asterisk) through the ACoA (white asterisk).

giography showed no aneurysm, no significant changes, and no other possible source of haemorrhage. The patient went on to make an uneventful and full recovery.

Discussion

Anomalies of the anterior cerebral-anterior communicating-recurrent artery complex are frequently encountered¹⁻⁵, especially during ACoA aneurysm surgery⁵. Among these anatomic variations, duplication of the proximal segment (A1) of the ACA is infrequent⁵. It has been reported to occur in 4% of subjects in cadaveric studies⁵ and in up to 0.5-9.7% of cases of ACoA aneurysm surgery^{1,4,6}. Although A1 segment duplication can be identified on angiography⁷, superimposition of vessels may render its identification difficult. Most clinically reported cases of A1 duplication are angiographically occult variations later identified during ACoA aneurysm surgery^{1,4,6} usually without consequences^{1,4,6,8,9}. In our case, the partially occluded A1 duplication was not recognized on preoperative CTA or angiogram as it mimicked an ACoA aneurysm. Thrombosis tends to occur at or adjacent to fenestrations and may explain the occlusion of one of the duplicated A1 segments in our patient¹⁰. Six previous cases of thrombosed intracranial arteries mimicking an aneurysm have been reported¹¹⁻¹⁵, including one case of a partially thrombosed fenestrated basilar artery that mimicked a vertebro-basilar junction aneurysm¹¹. All but two of these reported cases, led to surgical intervention¹¹⁻¹⁵. The decision to intervene surgically rather than embolize the supposed aneurysm was based on the risk of ACoA compromise in this patient whose anterior circulation of the right hemisphere arose exclusively from the ACoA. However, embolization of the partially occluded segment of the A1 duplication would not likely result in an ischemic event given that the occluded duplication segment is a blind pouch and no perforators were found to arise from it. Although no cause of SAH was identified, we do

not believe that the thrombosed duplicated vessel was responsible for the hemorrhage. The majority of blood was located around the ICA opposite to the A1 duplication and no parietal vessel abnormalities such as dissection or blister-like changes could be identified.

Post-operative review of the angiogram in this patient revealed the atypical location of the suspected aneurysm ipsilateral to the ICA occlusion. Given that aneurysms are known to form at sites of increased blood flow and hemodynamic stress, we should have expected this ACoA aneurysm to form on the left ACoA rather than on the right side, ipsilateral to the patient's ICA occlusion¹⁶⁻¹⁸. Asymmetry of A1 is associated with aneurysm formation and is found in 58-85% of patients harboring ACoA aneurysms¹⁹⁻²⁴. In these cases, aneurysm formation is three to five times more likely at the dominant than at the non-dominant A1-ACoA junction^{20,23} and up to four times more often at the dominant A-ACoA junction than at the ACoA itself²³. Furthermore, iatrogenic occlusion of the ICA results in an increased incidence of aneurysm formation at the contralateral side, most often at the ACoA¹⁶⁻¹⁸. To our knowledge, among the identified cases of ACoA aneurysm formation following iatrogenic occlusion of the ICA, there are no documented cases of an ACoA aneurysm forming truly ipsilateral to the occlusion²⁵⁻³⁹. Therefore, if the suspected image in our patient was in fact an aneurysm, one would have expected it to be found on the ACoA contralateral to the side of his ICA occlusion or broadly based on the ACoA itself. However, in our minds, this argument is not strong enough to prevent surgical intervention.

Conclusion

A1 segment duplication with one arm occluded, can mimic an ACoA aneurysm on angiography. The location of the anomaly at the ACoA on the side of least hemodynamic stress may provide a clue to suspect this variant.

References

- 1 Aydin IH, Takci E, Kadioglu HH, et al. Vascular variations associated with anterior communicating artery aneurysms: An intraoperative study. *Minim Invasive Neurosurg.* 1997; 40: 17-21.
- 2 Busse O. Aneurysmen und Bildungsfehler der Arteri communicans anterior. *Virchows Arch Pathol Anat.* 1921; 229: 178-189.
- 3 Maurer J, Maurer E, Perneczky A. Surgically verified variations in the A1 segment of the anterior cerebral artery: Report of two cases. *J Neurosurg.* 1991; 75: 950-953.
- 4 Ogawa A, Suzuki M, Sakurai Y, et al. Vascular anomalies associated with aneurysms of the anterior communicating artery: Microsurgical observations. *J Neurosurg.* 1990; 72: 706-709.
- 5 Perlmutter D, Rhoton AL Jr. Microsurgical anatomy of the anterior cerebral-anterior communicating-recurrent artery complex. *J Neurosurg.* 1976; 45: 259-272.
- 6 Suzuki M, Onuma T, Sakurai Y, et al. Aneurysms arising from the proximal (A1) segment of the anterior cerebral artery: Study of 38 cases. *J Neurosurg.* 1992; 76: 455-458.
- 7 Yamada T, Inagawa T, Takeda T. Ruptured aneurysm at the anterior cerebral artery fenestration. *J Neurosurg.* 1976; 45: 826-828.
- 8 Taylor R, Connolly ES, Duong H. Radiographic evidence and surgical confirmation of a saccular aneurysm on a hypoplastic duplicated A1 segment of the anterior cerebral artery: case report. *Neurosurgery.* 2000; 46(2): 482-484.
- 9 Friedlander RM, Oglivry CS. Aneurysmal subarachnoid hemorrhage in a patient with bilateral A1 fenestrations associated with an azygos anterior cerebral artery. Case report and literature review. *J Neurosurg.* 1996; 84 (4): 681-684.
- 10 Takahashi M, Kawanami H, Watanabe N, et al. Fenestration of the extracranial vertebral artery. *Radiology.* 1970; 96: 359-360.
- 11 Kalia KK, Pollack IF, Yonas H. A partially thrombosed, fenestrated basilar artery mimicking an aneurysm of the vertebrobasilar junction: case report. *Neurosurgery.* 1992; 30: 276-278.
- 12 Kawanishi M, Sakaguchi I, Miyake H. Occlusion of the posterior communicating artery mimicking cerebral aneurysm: case report. *Neurol Res.* 2003; 25: 543-545.
- 13 Komiyama M, Ishiguro T, Morikawa T, et al. Distal stump of an occluded intracranial vertebral artery at the vertebrobasilar junction mimicking a basilar artery aneurysm. *Acta Neurochir (Wien).* 2001; 143: 1013-1017.
- 14 Nakano S, Yokogami K, Ohta H, et al. A stump of occluded posterior cerebral artery mimicking a ruptured aneurysm: case report. *Int J Angiol.* 2000; 9: 51-52.
- 15 Pearl MS, Tamargo R, Gailloud P. Middle cerebral artery branch occlusion mimicking a saccular aneurysm on 3D digital subtraction angiography: Case report. *J Neurosurg.* 2008; 109: 1123-1125.
- 16 Drapkin AJ, Rose WS. Serial development of 'de novo' aneurysms after carotid ligation: case report. *Surg Neurol.* 1992; 38: 302-308.
- 17 Dyste GN, Beck DW. De novo aneurysm formation following carotid ligation: Case report and review of the literature. *Neurosurgery.* 1989; 24: 88-92.
- 18 Maiuri F, Spaziante R, Jacquetta G, et al. 'De novo' aneurysm formation: report of two cases. *Clin Neurol Neurosurg.* 1995; 97(3): 233-238.
- 19 Kirgis HD, Fisher WL, Llewellyn RC, et al. Aneurysms of the anterior communicating artery and gross anomalies of the circle of Willis. *J Neurosurg.* 1996; 25: 73-78.
- 20 Kwak R, Ohi T, Niizuma H, et al. Afferent artery and the site of neck of anterior communicating aneurysms. *Surg Neurol.* 1980; 13 (3): 221-223.
- 21 Rhoton AL. Anatomy of saccular aneurysms. *Surg Neurol.* 1980; 14: 59-66.
- 22 Stehbens WE. Aneurysms and anatomic variation of cerebral arteries. *Arch Pathol* 1963; 5: 45-64.
- 23 VanderArk GD, Kempe LC. Classification of anterior communication aneurysms as a basis for surgical approach. *J Neurosurg.* 1970; 32: 300-303.
- 24 Wilson G, Riggs HE, Rupp C. The pathologic anatomy of ruptured cerebral aneurysms. *J Neurosurg.* 1954; 11: 128-134.
- 25 Briganti F, Cirillo S, Caranci F, et al. Development of "de novo" aneurysms following endovascular procedures. *Neuroradiology.* 2002; 44 (7): 604-609.
- 26 Clark WC, Ray MW. Contralateral intracranial aneurysm formation as a late complication of carotid ligation. *Surg Neurol.* 1982; 18 (6): 458-462.
- 27 Fujiwara S, Fujii K, Fukui M. De novo aneurysm formation and aneurysm growth following therapeutic carotid occlusion for intracranial internal carotid artery (ICA) aneurysms. *Acta Neurochir (Wien).* 1993; 120 (1-2): 20-25.
- 28 German WJ, Black SP. Cervical ligation for internal carotid aneurysms. An extended follow-up. *J Neurosurg.* 1965; 23 (6): 572-577.
- 29 Heros RC. De Novo aneurysm formation following carotid ligation: case report and review of the literature. *Neurosurgery.* 1989; 24: 92.
- 30 Itoyama Y, Fujioka S, Takaki S, et al. Occlusion of internal carotid artery and formation of anterior communicating artery aneurysm in cervicocephalic fibromuscular dysplasia: follow-up case report. *Neurol Med Chir (Tokyo).* 1994; 34 (8): 547-550.
- 31 Jeffreys RV, Holmes AE. Common carotid ligation for the treatment of ruptured posterior communicating aneurysms. *J Neurol Neurosurg Psychiatry.* 1971; 34 (5): 576-579.
- 32 Matsuda M, Handa J, Saito A, et al. Ruptured cerebral aneurysms associated with arterial occlusion. *Surg Neurol.* 1983; 20: 4-12.
- 33 Miller JD, Jawad K, Jennett B. Safety of carotid ligation and its role in the management of intracranial aneurysms. *J Neurol Neurosurg Psychiatry.* 1977; 40 (1): 64-72.
- 34 Niino M, Shimozuru T, Nakamura K, et al. Long-term follow-up study of patients with cavernous sinus aneurysm treated by proximal occlusion. *Neurol Med Chir (Tokyo).* 2000; 40 (2): 88-96.
- 35 Shenkin HA, Polakoff P, Finneson BE. Intracranial internal carotid artery aneurysms; results of treatment by cervical carotid artery ligation. *J Neurosurg.* 1958; 15 (2): 183-191.
- 36 Somach FM, Shenkin HA. Angiographic end-results of carotid ligation in the treatment of carotid aneurysm. *J Neurosurg.* 1966; 24 (6): 966-974.
- 37 Timperman PE, Tomsick TA, Tew JM, et al. Aneurysm formation after carotid occlusion. *Am J Neuroradiol.* 1995; 16 (2): 329-331.
- 38 Wright RL, Sweet WH. Carotid or vertebral occlusion in the treatment of intracranial aneurysms; value of early and late readings of carotid and retinal pressures. *Clin Neurosurg.* 1963; 9: 163-192.
- 39 Salar G, Mingrino S. Development of intracranial saccular aneurysms: Report of two cases. *Neurosurgery.* 1981; 8: 462-465.

Michel W. Bojanowski, MD
 Department of Surgery, Division of Neurosurgery
 Centre Hospitalier de l'Université de Montreal (CHUM)
 Hôpital Notre-Dame - 1560 rue Sherbrooke Est
 Montreal, QC, Canada H2L 4M1
 Tel.: +1(514) 890-8000
 Fax: +1(514) 412-7816
 E-mail: michel.bojanowski.chum@ssss.gouv.qc.ca